

Cancer Immunology Miniatures

See related commentary by Lutz and Jaffee, p. 518

HLA-Binding Properties of Tumor Neoepitopes in Humans Edward F. Fritsch^{1,2,5}, Mohini Rajasagi^{1,2}, Patrick A. Ott^{2,4}, Vladimir Brusic^{1,4}, Nir Hacohen^{3,5}, and Catherine J. Wu^{1,2,4}**Abstract**

Cancer genome sequencing has enabled the rapid identification of the complete repertoire of coding sequence mutations within a patient's tumor and facilitated their use as personalized immunogens. Although a variety of techniques are available to assist in the selection of mutation-defined epitopes to be included within the tumor vaccine, the ability of the peptide to bind to patient MHC is a key gateway to peptide presentation. With advances in the accuracy of predictive algorithms for MHC class I binding, choosing epitopes on the basis of predicted affinity provides a rapid and unbiased approach to epitope prioritization. We show herein the retrospective application of a prediction algorithm to a large set of *bona fide* T cell–defined mutated human tumor antigens that induced immune responses, most of which were associated with tumor regression or long-term disease stability. The results support the application of this approach for epitope selection and reveal informative features of these naturally occurring epitopes to aid in epitope prioritization for use in tumor vaccines. *Cancer Immunol Res*; 2(6); 522–9. ©2014 AACR.

Introduction

We and others (1–3) have suggested that the vast number of personal, tumor-specific mutations found in the genome of patients with cancer provides a rich source of unique immunogens ("neoantigens") for use in tumor vaccination strategies. These tumor neoantigens are attractive as vaccine targets as they are expected to bypass the immune-dampening effects of central tolerance and because their expression is exquisitely tumor specific. To bring this highly personalized treatment approach to patients with cancer, one crucial challenge is the choice of which of the many possible personal mutated epitopes to incorporate in the vaccine.

Cancer genomes vary widely in the number of total and coding sequence mutations depending on the tumor type (4). The five most common tumor types in the United States (prostate, breast, lung, colon, and melanoma) harbor an average of 25 to 500 nonsynonymous coding sequence mutations. Vaccination approaches that use irradiated whole-tumor cells or various forms of cell lysates (5–11) have attempted to capture all such neoantigens (in addition to native tumor-associated antigens). Although these strategies seem comprehensive and have resulted in clinical benefit in some cases, they do not favor any particular T-cell immunogen. Hence, potentially highly effective immunogens may be

drowned out within the vast sea of immunologically irrelevant antigens. Such complete antigen preparations are similar to the endogenous presentation of the tumor cell to the immune system and lack the "pharmacologic specificity" of a rationally designed vaccine.

A more selective but still comprehensive approach for neoantigens could be envisioned by using every identified coding mutation as a separate immunogen. Although possibly feasible from the technical standpoint—especially for tumors with a low mutation load—the dilution of the potent immunogen is likely to reduce its effectiveness, and, thus, a more discriminating approach to identify the most effective subset seems advisable.

Potential Strategies to Identify and Prioritize Mutated Antigens

Multiple biochemical and biologic techniques are available that can help prioritize candidate mutated antigens for inclusion in tumor vaccines.

Mass spectrometry

Great strides in the fields of mass spectrometry (MS) and associated computational algorithms have enabled the characterization of the MHC-displayed "ligandome" (12, 13). This approach can be used to test whether a mutated peptide (or a native tumor-associated peptide) is displayed by tumor cells. This information is important as the peptide–MHC complexes are the substrates recognized by T-cell receptors (TCR). However, the approach is limited technically by insufficient amounts of tumor tissue and conceptually by the observation that few peptide-bound MHC targets are needed for an effective T-cell response (14, 15). As a result, many useful but less abundant targets on the cell may be bypassed in favor of those that are less potent but more highly represented.

Authors' Affiliations: ¹Cancer Vaccine Center; ²Department of Medical Oncology, Dana-Farber Cancer Institute; ³The Division of Allergy, Immunology, and Rheumatology, Department of Medicine, Massachusetts General Hospital; ⁴Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston; and ⁵Broad Institute of MIT and Harvard, Cambridge, Massachusetts

Corresponding Author: Catherine J. Wu, Dana-Farber Cancer Institute, Dana 540B, 450 Brookline Avenue, Boston, MA 02115. Phone: 617-632-5943; Fax: 617-632-6380; E-mail: cwu@partners.org

doi: 10.1158/2326-6066.CIR-13-0227

©2014 American Association for Cancer Research.

Ex vivo T-cell assays

Peripheral blood mononuclear cells or tumor-infiltrating lymphocytes can be tested in antigen-specific *ex vivo* assays to identify neoantigens that stimulate existing T-cell populations. This strategy would be expected to reveal the patient's natural response to neoantigens (2, 16). However, routine clinical application of *ex vivo* assays is costly and technically challenging, given the number of neoantigen mutations (requiring rapid preparation of many stimulatory immunogens), the requirement of MHC-matched antigen-presenting cells for some of these assays, and the relative insensitivity of these techniques. Most importantly, using *ex vivo* assays as a filter for neoantigen selection limits the spectrum of T-cell reactivity to existing T-cell responses. In patients with clinically evident tumors, this would restrict the selected neoantigen repertoire to the existing and possibly ineffective T-cell responses. It is currently unknown whether enhancement of an ongoing T-cell response or generation of *de novo* responses is clinically relevant for an effective tumor vaccine. Other biologic assays, such as *in vitro* or *in vivo* immunization of a humanized mouse, could also be considered, but such assays are likewise technically challenging, costly, and conceptually limited.

In silico prediction of peptide–MHC binding

Generation of an immune response to any mutated peptide sequence and recognition of tumor cells containing that peptide depend critically on the ability of the patient's MHC molecules to effectively bind to the mutated peptide and present it to a T cell. Advanced algorithms using neural network-based learning approaches have been developed to capitalize on large amounts of data describing peptides that bind with different strengths to a wide variety of class I MHC molecules (17). These algorithms allow rapid *in silico* prediction of peptide-binding strength to patient-specific MHC alleles, and potentially enable a more rapid and less restrictive approach to filter the list of candidate neoantigens from sequencing data. Using results from next-generation DNA sequencing, we have evaluated the binding for more than 100 different predicted peptides to understand the boundaries of the accuracy of prediction by these algorithms (18). To link this *in silico* analysis to potentially clinically and biologically relevant observations, we present here an analysis of 40 neoantigens previously identified as CD8⁺ T-cell targets in the literature.

The Predicted Binding Characteristics of Tumor Neoepitopes Recognized by T Cells in Patients with Antitumor Immunity

We have conducted an extensive search of the literature, including recent reviews on neoantigens (1, 2) from PubMed, and the most comprehensive list of cancer vaccine antigens compiled by the Cancer Research Institute (19), identifying reports of spontaneous CD8⁺ T-cell responses in patients with cancer in whom the target epitopes were discovered subsequently. To avoid bias of the results, reports of vaccinations with known epitopes or of selected searches for single T-cell epitopes (such as for an immune response to a known mutated oncogene) were not included. Multiple reports of spontaneous

CD8⁺ T-cell epitopes were identified, and, remarkably, in each case following an unbiased search for the dominant T-cell epitope, the target epitope was a neoantigen. Two thirds of the patients in these reports experienced significant partial or complete tumor regression or long-term stable disease, either spontaneously or following therapy.

As shown in Table 1, 31 of these 40 neoepitopes were identified in an unbiased manner based on cDNA expression cloning or MHC–peptide elution, while the remaining nine were found on the basis of genomic mutation and epitope-binding predictions. These neoantigens resulted from 35 missense mutations and five frameshift mutations (that led to novel open reading frames, neoORFs) and are restricted by 11 different HLA alleles, representing both common and less common alleles, as expected from sampling of the population at large. Approximately 80% of these are somatic mutations found exclusively in the tumors of individual patients. The remaining alterations are polymorphic loci within hematopoietically restricted minor histocompatibility antigens (miHAg) identified following hematopoietic stem cell transplantation for blood malignancies. In almost every case, the mutated peptide was significantly (>100×) more potent than the cognate native peptide in the induction of T-cell IFN γ production or cytotoxicity. These examples represent seven different cancer types (non–small cell lung cancer, melanoma, renal cell carcinoma, bladder cancer, B-cell acute lymphoblastic leukemia, multiple myeloma, and chronic lymphocytic leukemia).

Because these neoepitopes are associated with biologic responses, they provide an ideal set of sequences for retrospective peptide affinity predictions to "reverse engineer" predictable characteristics of effective epitopes. For this analysis, we used the netMHCpanv2.4 algorithm (Center for Biological Sequence Analysis, Technical University of Denmark, Lyngby, Denmark; www.dtu.dk; ref. 20). NetMHCpan is an artificial neural network–trained algorithm with an extensive training dataset (17), including 43 HLA-A and HLA-B alleles, representing approximately 90% and 60%, respectively, of the allelic population distribution, with more than 1,000 members each in the training set. NetMHCpan was determined to be one of the most accurate predictive algorithms in a 2012 competition (21). We applied this algorithm to individually predict MHC binding for all possible tiled peptides containing the mutated or the corresponding unmutated residues of these observed spontaneous epitopes to determine:

- whether the naturally recognized epitopes would have been predicted;
- the predicted affinity of each mutated epitope; and
- the predicted affinity of each cognate native epitope (focusing on only the missense mutations).

Functional neoepitopes are correctly predicted by a class I MHC–peptide binding algorithm

Thirty-one of the epitopes shown in Table 1 were identified by *ex vivo* T-cell reactivity or MS and did not use genomic sequence or binding prediction information as a component of their identification. For all but one of these 31 epitopes, we found that the reported epitope was the peptide with the

Table 1. Biologic features and predicted binding affinities of neoantigen-directed T-cell responses in humans

Group	Gene (Ref.)	Identification approach ^a	Favorable clinical response	MUT>>>NAT T-cell response detected (approach) ^b	HLA allele	Mutated epitope (native allele)	Predicted binding affinity (IC ₅₀ nmol/L)	
							MUT	NAT
1	ECGF-1 (ref. 32; miHAg)	cDNA	Yes	Yes (Y)	B*07:02	RPHAIRRPLAL (R)	3	2
	ME-1 (31)	cDNA	Yes	Yes (C)	A*02:01	FLDEFMEGV (A)	3	2
	PLEKHM2 (16)	WES	Yes	Yes (Y)	A*01:01	LTDDRLLFTCY (H)	3	97
	FNDC3B (18)	WES	Yes	±(T) (≥10×)	A*02:01	VWMSWAPPV (L)	4	7
	PRDX5 (27)	cDNA	NR	Yes (C)	A*02:01	LLLDLLVSI (S)	5	7
	MATN2 (16)	WES	Yes	Yes (Y)	A*11:01	KTLTTSVFK (E)	5	20
	DDX21 (33)	cDNA	Yes	Yes (C)	A*68:01	EAFIQPIR (S)	10	29
	RBAF (29)	cDNA	Yes	Yes (C)	B*07:02	RPHVPESAF (G)	10	68
	GAS7 (16, 34)	cDNA (and later WES)	Yes	Yes (C)	A*02:01	SLADEAEVYL (H)	12	39
	ATR (35)	WES	Yes	±(T) (≥10×)	A*03:01	KLYEPLLK (S)	13	13
	SIRT2 (29)	cDNA	Yes	±C (10×)	A*03:01	KIFSEVTLK (P)	14	16
	EF2 (36)	MS	NR	Yes (C)	A*68:02	ETVSEGSNV (E)	16	27
	KIAA0223 (HA-1; ref. 37; miHAg)	MS	Yes (severe GVHD)	NR	A*02:01	VLHDDLLEA (R)	17	140
	GAPDH (34)	cDNA	Yes	Yes (C)	A*02:01	GIVEGLITV (M)	21	27
	BCL2A1 (ref. 38; miHAg) ^a	Linkage	NR	Yes (C)	A*24:02	DYLQYVLQI (C)	22	34
	HSP70 (39)	cDNA	NR	Yes (C)	A*02:01	SLFEGDIYT (F)	23	7
	ACTININ (30)	cDNA	Yes	Yes (C)	A*02:01	FIASNGVKLV (K)	29	44
	CDK12 (16)	WES	Yes	Yes (Y)	A*11:01	CILGKLFTE (E)	33	42
	KIAA1440 (40)	cDNA	Yes	Yes (C)	A*01:01	QTACEVLDY (T)	33	78
	HAUS3 (16)	WES	Yes	Yes (Y)	A*02:01	ILNAMIAKI (T)	34	36
	BCL2A1 (ref. 38; miHAg) ^a	Linkage	NR	Yes (C)	B*44:03	KEFEDDIINW (G)	36	27
	PPP1R3B (16, 28)	cDNA (and later WES)	Yes	Yes (Y)	A*01:01	YTDFHCQYV (P)	49	72
	HB-1 (ref. 41; miHAg)	cDNA	NR	Yes (C)	B*44:03	EEKRGSLLHWV (Y)	81	67
MUM-2 (42)	cDNA	Yes	Yes (C)	B*44:02	SELFRRSGLDSY (R)	184	182	
KIAA0205 (43)	cDNA	NR	Yes (C)	B*44:03	AEPIDIQITW (N)	258	288	
GPMMB (29)	cDNA	Yes	Yes (C)	A*03:01	TLDWLLQTPK (G)	282	179	
CSNK1A1 (16)	WES	Yes	Yes (Y)	A*02:01	GLFGDIYLA (S)	6	1,312	
CLPP (44)	cDNA	Yes	Yes (C)	A*02:01	ILDKVLVHL (P)	32	7,566	
CTNMB1 (45)	cDNA	Yes (?)	Yes (C)	A*24:02	SYLDSGIHF (S)	41	18,746	
SNRP116 (29)	cDNA	Yes	Yes (C)	A*03:01	KILDAWAQQK (E)	48	14,976	
OS9 (46)	cDNA	NR	Yes (C)	B*44:03	KELEGILLI (P)	60	1,161	

(Continued on the following page)

Table 1. Biologic features and predicted binding affinities of neoantigen-directed T-cell responses in humans (Cont'd)

Group	Gene (Ref.)	Identification approach ^a	Favorable clinical response	MUT>>>NAT T-cell response detected (approach) ^b	HLA allele	Mutated epitope (native allele)	Predicted binding affinity (IC ₅₀ nmol/L)	
							MUT	NAT
3	<i>MYH2</i> (47)	cDNA	Yes	Yes (C)	A*03:01	K INKNPKYK (E)	141	4,960
	<i>MART-2</i> (48)	cDNA	Yes (weak)	Yes (C)	A*01:01	F L EGNEVGKTY (G)	1,115	4,504
	<i>NFYC</i> (49)	cDNA	NR	Yes (C)	B*52:01	AQQITKTEV (Q)	7,314	5,701
neoORF	<i>CDK4</i> (50)	cDNA	NR	Yes (C)	A*02:01	ACDPHSGHFV (R)	11,192	25,222
	<i>pARF14-ORF3</i> (51)	cDNA	Yes	Not relevant	A*11:01	AVCPWTWLR	25	Not relevant
	<i>HMSD-γ</i> (ref. 52; miHAg)	cDNA	Yes	Not relevant	B*44:03	MEFIEVFSHF	36	Not relevant
	<i>PANE-1</i> (ref. 53; miHAg)	MS	NR	Not relevant	A*03:01	RWVDLPGVLK	44	Not relevant
	<i>MUM1</i> (54)	cDNA	Yes	Yes (C)	B*44:02	(L)EEKLVVLF ^c (S)	434 (409)	Not relevant
	<i>P2×5</i> (ref. 55; miHAg)	Linkage	Yes	Not relevant	B*07:02	TPNQRQNVK	1,769	Not relevant

NOTE: Predicted affinity highlighting: green type, strong/moderate binder (IC₅₀ ≤ 150 nmol/L); blue type, weak binder (IC₅₀ between 150 and 500 nmol/L); and black type, nonbinder (IC₅₀ ≥ 500 nmol/L).

Bold, identification by binding predictions and either genomic sequence or genetic linkage.

Underlined values: comparable experimentally measured binding affinity of mutated and native peptides were reported.

Abbreviations: cDNA, cDNA library expression cloning; NR, not reported; WES, whole-exome sequencing.

^aTwo separate CTL clones specific for peptides restricted by distinct MHC alleles (A2402 and B4403) were identified. Each peptide spanned a distinct missense mutation in the *BCL2A1* gene.

^bC, cytotoxicity; γ, IFN_γ stimulation; T, CD107a stimulation.

^cSpecial case of a missense, partial neoORF.

strongest predicted MHC-binding affinity among the tiled peptides containing the mutation. The only exception was a *MUMI*-derived 10mer containing an additional leucine at the N-terminus that had a slightly better predicted affinity (IC_{50} , 409 nmol/L) than the observed 9mer (IC_{50} , 434 nmol/L). We conclude that the MHC-peptide binding prediction algorithm netMHCpan consistently predicts the naturally recognized tumor neoepitope from all of the possible epitopes harboring a specific mutation.

Most functional neoepitopes have strong to moderate predicted IC_{50}

Twenty of 31 (65%) of the naturally recognized missense and neoORF epitopes had predicted $IC_{50} < 50$ nmol/L (strong binders) and three of 31 (10%) had a predicted IC_{50} between 50 and 150 nmol/L (moderate binders). Thus, 75% of the dominant T-cell clones isolated from the naturally occurring T-cell populations recognize an epitope with a strong or moderate predicted affinity ($IC_{50} < 150$ nmol/L) for the patient's MHC allele. Because an unbiased functional assay (cytolysis, $IFN\gamma$ production, or MS) was the critical test used to identify each of the stimulating peptides in these 31 examples, it is unlikely that there was an experimental bias toward the identification of epitopes with higher predicted affinity. Four of 31 naturally recognized peptides were predicted to be "weak" binding peptides (IC_{50} between 150 and 500 nmol/L), indicating that a total of 27 of 31 (87%) of the naturally occurring epitopes would have been considered as binding peptides ($IC_{50} < 500$ nmol/L affinity) using netMHCpan.

Conversely, only four of 31 naturally recognized peptides were predicted by netMHCpan to be nonbinders ($IC_{50} > 500$ nmol/L); they may be false negatives from the prediction algorithm or may represent low affinity yet functional epitopes. Although these alternatives cannot be distinguished on the basis of the available data, three observations are relevant. First, for the three epitopes arising from missense mutations (*MART-2*, *NFYC*, and *CDK4*), cytolytic activity was preferentially induced by the mutated peptide and not by the native peptide at a range of peptide concentrations (1–10 nmol/L) comparable with those observed with more strongly predicted binding peptides. Second, for the Arg \rightarrow Cys *CDK4* mutation, the highly oxidizable sulfur residue may contribute serendipitously to MHC binding as a "pseudo-" anchor residue that could not have been accounted for by the prediction algorithms. Finally, T cells recognizing the fourth epitope (the miHAg *P2* \times 5) represented as much as 1.6% of all circulating T cells following the therapeutic infusion of donor lymphocytes. Results from these 40 examples dataset suggest that there are limitations to the capability of predictive algorithms and that up to 15% of target T-cell epitopes may be missed by the prediction algorithms.

Most of the cognate native peptides are predicted to bind MHC equally to the mutated peptides

In addition to analyzing MHC binding to the mutated epitopes, we also compared the predicted affinities of the cognate native epitopes corresponding to all 35 missense epitopes in Table 1 and identified three distinct classes. The

predominant class (26 of 35 or 74%; group 1) was composed of native/mutated pairs that were predicted to bind with comparable affinity [with 23 of 26 showing strong to moderate predicted binding ($IC_{50} < 150$ nmol/L) and the remaining three showing weak binding (IC_{50} between 150 and 500 nmol/L)]. Despite comparable predicted binding, in almost all cases, the mutated peptide had been found to be significantly more potent in stimulating T cells than the native peptide. A smaller group (6 of 35 or 17%; group 2) showed low predicted binding for the native epitope and strong binding for the mutated epitope, directly correlating with the differential T-cell responses to the mutated and native peptides. Finally, in a minority of cases both mutated and native peptides were predicted to be nonbinding (3 of 35 or 9%; group 3). We note that each group in Table 1 comprises multiple HLA with no apparent bias in representation. Although the existence of the group 1 and 2 epitopes is not surprising, the predominance of the group 1 epitopes (containing 74% of all missense epitopes and four times more abundant than group 2) with comparable affinities for both the mutated and native peptides is unexpected.

Discussion

The MHC-bound peptide can be considered as a double-sided "key," which must fit both the MHC and the TCR "locks" to stimulate an immune response and for subsequent target-cell cytolysis (Fig. 1A). Sequence-specific binding of peptides to the MHC molecule is highly dependent on the interactions of the peptide side chains at particular positions ("anchors") along the length of the peptide with chemical moieties defined by the polymorphic residues that constitute the MHC-binding pocket (22–24); hence, predictive calculations are sequence dependent (25). Furthermore, analysis of these critical MHC-binding positions and residues over a wide range of MHC alleles shows that only a few positions of the peptide are anchor positions and only a few amino acids at the anchor positions of the peptide contribute to binding in a positive manner (26). On the other side of the "key," TCR recognition of the peptide-MHC complex gains specificity from the ordered presentation of the other face of the peptide conferred by the anchoring residues.

For the majority of the missense mutations, both the native and the mutated peptides were predicted to be binding peptides (group 1; Fig. 1B). This observation is almost certainly a consequence of the mutations affecting the region of the peptide "key" that is involved in TCR recognition. In all but two of these 26 examples, the mutation was in a nonanchor position (as identified by the online tool provided at <http://www.syfpeithi.de/>; ref. 26). In the two nonconforming examples (*PLEKHM2* and *KIAA1440*), a second anchor residue was already present in the native peptide. Other investigators have also reported mutations with equivalent affinity predictions for the native and the mutated peptide pairs (16, 27). Our broader analysis suggests that such mutant epitopes are a common phenomenon. Only a minority of the missense mutations were found in group 2 characterized by nonbinding of the native peptide. All of the group 2 examples, except for *MYOSIN*, were




Figure 1. A, the two faces of a bound peptide to the MHC and TCR molecules form a "double-sided key" that must be present to stimulate an antigen-specific immune response. Green, anchor residues in the peptide that interact with MHC. Purple, regions of the peptide that interact with the TCR surface. B, a scatterplot of the predicted affinities of epitopes that stimulate detectable neoantigen T-cell responses, shown in Table 1. Group 1 epitopes demonstrate comparable predicted affinities of native and mutated peptides and were determined to have mutations in regions of the peptide critical for interactions with the TCR (dark purple, strong/moderate binders; light purple, weak binders). Group 2 epitopes (green) are mutated peptides with strong/moderate predicted affinity whose corresponding native peptides are not predicted to bind MHC, and were found to have mutations in the peptide residues critical for the interaction with MHC. Group 3 epitopes (gray) represent peptides in which neither the native nor mutated peptide are predicted to be HLA-binding peptides and may be either false negatives of the prediction algorithm or very low-affinity functional epitopes.

mutations to preferred anchor residues at critical anchor positions.

Although the majority of the naturally occurring tumor epitopes were derived from the corresponding native peptides predicted to bind MHC, the vast majority (>98%) of the native human peptidome is not predicted to contain peptides that are binding epitopes of human MHC (our unpublished analysis using netMHCpan). Random mutational events that convert a nonbinding peptide (the vastly predominant target) to a binding peptide (group 2) are expected to be rare because they require mutation to one of only a few specific amino acids at a small number of anchor positions. For most MHC molecules, there are only one or two important anchor positions, and usually only two or three amino acids at those positions promote binding. Conversely, nonanchor positions are three to four times more abundant than anchor positions, and most mutations to native binding epitopes in these nonanchor positions would maintain MHC binding (group 1). This simple probabilistic explanation may be sufficient to account for the predominance of the observed group 1 epitopes (derived from the vastly underrepresented class of native peptides that are predicted to bind MHC). Alternatively, more complex explanations may be required. For example, aspects of central immune tolerance that are currently not well understood may cause the extant TCR repertoire to more effectively respond to peptides presenting a surface chemically distinct from any native peptide (group 1) than to peptides that more efficiently present an otherwise native peptide surface (group 2).

We did not observe weak native peptide binders that converted to strong/moderate mutated binders or strong/moderate native peptide binders that converted to weak mutated peptide binders. Although this may reflect the relatively limited dataset we used, it could also be that such upgrading or downgrading of binding involves anchor residue changes that moderately increase/decrease binding affinity but retain similar chemical structure of the nonanchor residues available for TCR recognition. In these scenarios, because the native peptides could bind to MHC, central immune tolerance may have effectively deleted cells with the reactive TCR, rendering both native and mutated peptides nonimmunogenic.

From the perspective of vaccine efficacy, we propose that mutations resulting in either group 1 or 2 binders should be considered as acceptable for use as immunogens, as both types of mutated neoepitopes have been found *in vivo* in patients with cancer with spontaneous tumor regressions and in long-term cancer survivors. Notably, in long-term survivors, T cells specific to mutated tumor epitopes from both groups 1 and 2 have been found to persist over many years (28, 29).

From the perspective of safety, there have been no reports of immune-mediated toxicities (except for the expected occurrence of GVHD as a result of responses against miHAGs) despite the observation that for most of the mutations, the cognate native peptide was predicted and experimentally demonstrated in some cases (18, 27, 30, 31), to bind MHC as well as the mutated peptide (group 1). Importantly, in almost all cases the mutant peptide was shown to be more potent than

the native peptide in stimulating T-cell cytotoxicity or IFN γ production. The absence of autoimmune toxicity in these patients fits with the model that T cells reactive to the native epitope were eliminated by central immune tolerance and that T cells reactive to the mutated epitope do not cross-react to the native epitope as the mutation exclusively affects the TCR binding region.

In conclusion, *in silico* peptide-binding predictions provide a useful and rapidly deployable tool to capture the types of immunogens that are naturally observed in patients with cancer, many of whom experienced tumor regression and sometimes long-term tumor control. Moreover, results from our retrospective prediction study reveal features of these epitopes to further guide inclusion as immunogens in vaccines. We have recently initiated a clinical study using personalized neoantigen epitopes identified by whole-exome sequencing and prioritized by MHC-binding predictions in which we will carefully monitor the immune response to each mutation (NCT01970358).

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Hacohen N, Fritsch EF, Carter TA, Lander ES, Wu C. Getting personal with neoantigen-based therapeutic cancer vaccines. *Cancer Immunol Res* 2013;1:11–15.
- Heemskerk B, Kvistborg P, Schumacher TN. The cancer antigenome. *EMBO J* 2013;32:194–203.
- Castle JC, Kreiter S, Diekmann J, Lower M, van der Roemer N, de Graaf J, et al. Exploiting the mutanome for tumor vaccination. *Cancer Res* 2012;72:1081–91.
- Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 2013;499:214–8.
- Berd D, Murphy G, Maguire HC Jr, Mastrangelo MJ. Immunization with haptenized, autologous tumor cells induces inflammation of human melanoma metastases. *Cancer Res* 1991;51:2731–4.
- Soiffer R, Hodi FS, Haluska F, Jung K, Gillessen S, Singer S, et al. Vaccination with irradiated, autologous melanoma cells engineered to secrete granulocyte-macrophage colony-stimulating factor by adenoviral-mediated gene transfer augments antitumor immunity in patients with metastatic melanoma. *J Clin Oncol* 2003;21:3343–50.
- Randazzo M, Terness P, Opelz G, Kleist C. Active-specific immunotherapy of human cancers with the heat shock protein Gp96-revisited. *Int J Cancer* 2012;130:2219–31.
- Senzer NN, Kaufman HL, Amatruda T, Nemunaitis M, Reid T, Daniels S, et al. Phase II clinical trial of a granulocyte-macrophage colony-stimulating factor-encoding, second-generation oncolytic herpesvirus in patients with unresectable metastatic melanoma. *J Clin Oncol* 2009;27:5763–71.
- Morton DL, Foshag LJ, Hoon DS, Nizze JA, Famatiga E, Wanek LA, et al. Prolongation of survival in metastatic melanoma after active specific immunotherapy with a new polyvalent melanoma vaccine. *Ann Surg* 1992;216:463–82.
- Vaishampayan U, Abrams J, Darrah D, Jones V, Mitchell MS. Active immunotherapy of metastatic melanoma with allogeneic melanoma lysates and interferon alpha. *Clin Cancer Res* 2002;8:3696–701.
- Bystryn JC, Zeleniuch-Jacquotte A, Oratz R, Shapiro RL, Harris MN, Roses DF. Double-blind trial of a polyvalent, shed-antigen, melanoma vaccine. *Clin Cancer Res* 2001;7:1882–7.
- Rammensee HG, Singh-Jasuja H. HLA ligandome tumor antigen discovery for personalized vaccine approach. *Expert Rev Vaccines* 2013;12:1211–7.
- Walter S, Weinschenk T, Stenzl A, Zdrojowy R, Pluzanska A, Szczylik C, et al. Multipeptide immune response to cancer vaccine IMA901 after single-dose cyclophosphamide associates with longer patient survival. *Nat Med* 2012;18:1254–61.
- Purbhoo MA, Irvine DJ, Huppa JB, Davis MM. T cell killing does not require the formation of a stable mature immunological synapse. *Nat Immunol* 2004;5:524–30.
- Sykulev Y, Joo M, Vturina I, Tsomides TJ, Eisen HN. Evidence that a single peptide–MHC complex on a target cell can elicit a cytolytic T cell response. *Immunity* 1996;4:565–71.
- Robbins PF, Lu YC, El-Gamil M, Li YF, Gross C, Gartner J, et al. Mining exomic sequencing data to identify mutated antigens recognized by adoptively transferred tumor-reactive T cells. *Nat Med* 2013;19:747–52.
- Lundegaard C, Lund O, Nielsen M. Prediction of epitopes using neural network based methods. *J Immunol Methods* 2011;374:26–34.
- Rajasagi M, Keskin D, Kim J, Zhang W, Shukla S, DeLuca D, et al. Systematic identification of personal mutated tumor-specific neoantigens in CLL [abstract]. In: Proceedings of the 54th ASH Annual Meeting and Exposition; 2012 Dec 8–11. Atlanta, GA: Georgia World Congress Center. Abstract nr 954. [cited 2013 August]. Available from: <https://ash.confex.com/ash/2012/webprogram/Paper51216.html>.
- van der Bruggen P, Stroobant V, Vigneron N, Van den Eynde BJ. Peptide database: T cell-defined tumor antigens resulting from mutations. Available from: <http://cancerimmunity.org/peptide/mutations/>.
- Center for Biological Sequence Analysis. Lyngby, Denmark: Technical University of Denmark. Available from: www.dtu.dk.
- Zhang GL, Ansari HR, Bradley P, Cawley GC, Hertz H, Hu X, et al. Machine learning competition in immunology—prediction of HLA class I binding peptides. *J Immunol Methods* 2011;374:1–4.
- Yanover C, Bradley P. Large-scale characterization of peptide–MHC binding landscapes with structural simulations. *Proc Natl Acad Sci U S A* 2011;108:6981–6.
- Matsumura M, Fremont DH, Peterson PA, Wilson IA. Emerging principles for the recognition of peptide antigens by MHC class I molecules. *Science* 1992;257:927–34.
- Fremont DH, Matsumura M, Stura EA, Peterson PA, Wilson IA. Crystal structures of two viral peptides in complex with murine MHC class I H-2Kb. *Science* 1992;257:919–27.

Authors' Contributions

Conception and design: E.F. Fritsch, M. Rajasagi, N. Hacohen, C.J. Wu
Development of methodology: E.F. Fritsch, M. Rajasagi
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): E.F. Fritsch, C.J. Wu
Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): E.F. Fritsch, V. Brusic, N. Hacohen
Writing, review, and/or revision of the manuscript: E.F. Fritsch, P.A. Ott, V. Brusic, N. Hacohen, C.J. Wu
Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): E.F. Fritsch, C.J. Wu
Study supervision: N. Hacohen, C.J. Wu

Acknowledgments

The authors thank Sachet Shukla for helpful and insightful discussions.

Grant Support

The work on neoepitope-based vaccines was financially supported by the Blavatnik Family Foundation and the NIH (NHLBI:5 R01 HL103532-03; NCI:1R01CA155010-02).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received December 28, 2013; revised February 5, 2014; accepted February 17, 2014; published OnlineFirst March 3, 2014.

25. Sidney J, Assarsson E, Moore C, Ngo S, Pinilla C, Sette A, et al. Quantitative peptide binding motifs for 19 human and mouse MHC class I molecules derived using positional scanning combinatorial peptide libraries. *Immunome Res* 2008;4:2.
26. Rammensee H, Bachmann J, Emmerich NP, Bachor OA, Stevanovic S. SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 1999;50:213–9.
27. Sensi M, Nicolini G, Zanon M, Colombo C, Molla A, Bersani I, et al. Immunogenicity without immunoselection: a mutant but functional antioxidant enzyme retained in a human metastatic melanoma and targeted by CD8(+) T cells with a memory phenotype. *Cancer Res* 2005;65:632–40.
28. Lu YC, Yao X, Li YF, El-Gamil M, Dudley ME, Yang JC, et al. Mutated PPP1R3B is recognized by T cells used to treat a melanoma patient who experienced a durable complete tumor regression. *J Immunol* 2013;190:6034–42.
29. Lennerz V, Fatho M, Gentilini C, Frye RA, Lifke A, Ferel D, et al. The response of autologous T cells to a human melanoma is dominated by mutated neoantigens. *Proc Natl Acad Sci U S A* 2005;102:16013–8.
30. Echchakir H, Mami-Chouaib F, Vergnon I, Baurain JF, Karanikas V, Chousib S, et al. A point mutation in the alpha-actinin-4 gene generates an antigenic peptide recognized by autologous cytolytic T lymphocytes on a human lung carcinoma. *Cancer Res* 2001;61:4078–83.
31. Karanikas V, Colau D, Baurain JF, Chiari R, Thonnard J, Gutierrez-Roelens I, et al. High frequency of cytolytic T lymphocytes directed against a tumor-specific mutated antigen detectable with HLA tetramers in the blood of a lung carcinoma patient with long survival. *Cancer Res* 2001;61:3718–24.
32. Slager EH, Honders MW, van der Meijden ED, van Luxemburg-Heija SA, Kloosterboer FM, Kester MG, et al. Identification of the angiogenic endothelial-cell growth factor-1/thymidine phosphorylase as a potential target for immunotherapy of cancer. *Blood* 2006;107:4954–60.
33. Baurain JF, Colau D, van Baren N, Landry C, Martelange V, Vikkula M, et al. High frequency of autologous anti-melanoma CTL directed against an antigen generated by a point mutation in a new helicase gene. *J Immunol* 2000;164:6057–66.
34. Zhou J, Dudley ME, Rosenberg SA, Robbins PF. Persistence of multiple tumor-specific T-cell clones is associated with complete tumor regression in a melanoma patient receiving adoptive cell transfer therapy. *J Immunother* 2005;28:53–62.
35. van Rooij N, van Buuren MM, Philips D, Velds A, Toebes M, Heemskerk B, et al. Tumor exome analysis reveals neoantigen-specific T-cell reactivity in an ipilimumab-responsive melanoma. *J Clin Oncol* 2013;31:e439–42.
36. Hogan KT, Eisinger DP, Cupp SB III, Lekstrom KJ, Deacon DD, Shabanowitz J, et al. The peptide recognized by HLA-A68.2-restricted, squamous cell carcinoma of the lung-specific cytotoxic T lymphocytes is derived from a mutated elongation factor 2 gene. *Cancer Res* 1998;58:5144–50.
37. den Haan JM, Meadows LM, Wang W, Pool J, Blokland E, Bishop TL, et al. The minor histocompatibility antigen HA-1: a diallelic gene with a single amino acid polymorphism. *Science* 1998;279:1054–7.
38. Akatsuka Y, Nishida T, Kondo E, Miyazaki M, Tajiri H, Lida H, et al. Identification of a polymorphic gene, BCL2A1, encoding two novel hematopoietic lineage-specific minor histocompatibility antigens. *J Exp Med* 2003;197:1489–500.
39. Gaudin C, Kremer F, Angevin E, Scott V, Triebel F. A hsp70-2 mutation recognized by CTL on a human renal cell carcinoma. *J Immunol* 1999;162:1730–8.
40. Zhou X, Jun DY, Thomas AM, Huang X, Huang LQ, Mautner J, et al. Diverse CD8⁺ T-cell responses to renal cell carcinoma antigens in patients treated with an autologous granulocyte-macrophage colony-stimulating factor gene-transduced renal tumor cell vaccine. *Cancer Res* 2005;65:1079–88.
41. Dolstra H, Fredrix H, Maas F, Coulie PG, Brasseur F, Mensink E, et al. A human minor histocompatibility antigen specific for B cell acute lymphoblastic leukemia. *J Exp Med* 1999;189:301–8.
42. Chiari R, Foury F, De Plaen E, Baurain JF, Thonnard J, Coulie PG. Two antigens recognized by autologous cytolytic T lymphocytes on a melanoma result from a single point mutation in an essential house-keeping gene. *Cancer Res* 1999;59:5785–92.
43. Gueguen M, Patard JJ, Gaugler B, Brasseur F, Renaud JC, Van Cangh PJ, et al. An antigen recognized by autologous CTLs on a human bladder carcinoma. *J Immunol* 1998;160:6188–94.
44. Corbiere V, Chapiro J, Stroobant V, Ma W, Lurquin C, Lethe B, et al. Antigen spreading contributes to MAGE vaccination-induced regression of melanoma metastases. *Cancer Res* 2011;71:1253–62.
45. Robbins PF, El-Gamil M, Li YF, Kawakami Y, Loftus D, Appella E, et al. A mutated beta-catenin gene encodes a melanoma-specific antigen recognized by tumor infiltrating lymphocytes. *J Exp Med* 1996;183:1185–92.
46. Vigneron N, Ooms A, Morel S, Degiovanni G, Van Den Eynde BJ. Identification of a new peptide recognized by autologous cytolytic T lymphocytes on a human melanoma. *Cancer Immunol* 2002;2:9.
47. Zorn E, Hercend T. A natural cytotoxic T cell response in a spontaneously regressing human melanoma targets a neoantigen resulting from a somatic point mutation. *Eur J Immunol* 1999;29:592–601.
48. Kawakami Y, Wang X, Shofuda T, Sumimoto H, Tupesis J, Fitzgerald E, et al. Isolation of a new melanoma antigen, MART-2, containing a mutated epitope recognized by autologous tumor-infiltrating T lymphocytes. *J Immunol* 2001;166:2871–7.
49. Takenoyama M, Baurain JF, Yasuda M, So T, Sugaya M, Hanagiri T, et al. A point mutation in the NFYC gene generates an antigenic peptide recognized by autologous cytolytic T lymphocytes on a human squamous cell lung carcinoma. *Int J Cancer* 2006;118:1992–7.
50. Wölfel T, Hauer M, Schneider J, Serrano M, Wolfel C, Klehmann-Hieb E, et al. A p16INK4a-insensitive CDK4 mutant targeted by cytolytic T lymphocytes in a human melanoma. *Science* 1995;269:1281–4.
51. Huang J, El-Gamil M, Dudley ME, Li YF, Rosenberg SA, Robbins PF. T cells associated with tumor regression recognize frameshifted products of the CDKN2A tumor suppressor gene locus and a mutated HLA class I gene product. *J Immunol* 2004;172:6057–64.
52. Kawase T, Akatsuka Y, Torikai H, Morishima S, Oka A, Tsujimura A, et al. Alternative splicing due to an intronic SNP in HMSD generates a novel minor histocompatibility antigen. *Blood* 2007;110:1055–63.
53. Brickner AG, Evans AM, Mito JK, Xuereb SM, Feng X, Nishida T, et al. The PANE1 gene encodes a novel human minor histocompatibility antigen that is selectively expressed in B-lymphoid cells and B-CLL. *Blood* 2006;107:3779–86.
54. Coulie PG, Lehmann F, Lethe B, Herman J, Luiguin C, Andrawiss M, et al. A mutated intron sequence codes for an antigenic peptide recognized by cytolytic T lymphocytes on a human melanoma. *Proc Natl Acad Sci U S A* 1995;92:7976–80.
55. de Rijke B, van Horsen-Zoetbrood A, Beekman JM, Otterud B, Maas F, Woestenenk R, et al. A frameshift polymorphism in P2 × 5 elicits an allogeneic cytotoxic T lymphocyte response associated with remission of chronic myeloid leukemia. *J Clin Invest* 2005;115:3506–16.

Cancer Immunology Research

HLA-Binding Properties of Tumor Neoepitopes in Humans

Edward F. Fritsch, Mohini Rajasagi, Patrick A. Ott, et al.

Cancer Immunol Res 2014;2:522-529. Published OnlineFirst March 3, 2014.

Updated version Access the most recent version of this article at:
doi:[10.1158/2326-6066.CIR-13-0227](https://doi.org/10.1158/2326-6066.CIR-13-0227)

Cited articles This article cites 52 articles, 35 of which you can access for free at:
<http://cancerimmunolres.aacrjournals.org/content/2/6/522.full#ref-list-1>

Citing articles This article has been cited by 26 HighWire-hosted articles. Access the articles at:
<http://cancerimmunolres.aacrjournals.org/content/2/6/522.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerimmunolres.aacrjournals.org/content/2/6/522>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.